

# AUTHOR INDEX

- Abrams H, A-16 (No. 3)  
Acute Respiratory Distress Syndrome Network, 16  
Agnelli G, 55  
Aickin MI, 36  
Alberti C, 93  
Alberts DS, 21  
Alberts MJ, 8  
Anand SS, 3  
Anderson DC, 55  
Ankum, WM, 33  
Appleton S, 97  
Armstrong DG, 107  
Arroll B, 14  
Asplund K, 8  
Attia J, A-11-12 (No. 3)  
Atwood CW, 45
- Badger CM, 56  
Balden E, 85  
Bao B, 59  
Barbui C, 84  
Bastyr EJ 3d, 88  
Bates ER, 4  
Beasley R, 32  
Beck FW, 59  
Beck RW, 81  
Becker DM, 62, 82  
Belda J, 97  
Bellomo R, 41  
Bender BS, 90  
Bender R, 17  
Berger M, 17  
Berkus MD, 87  
Berlin JA, 77  
Beyth RJ, 54  
Bhatt DL, 49  
Black S, 117  
Bloch R, 74-75  
Boaz M, 91  
Bond JH, 103  
Bonithon-Kopp C, 104  
Borenstein P, 111  
Borwick A, 57  
Brain Attack Coalition, 8  
Brenner BE, 46  
Bridges CB, 90  
Brochard L, 16  
Brodows RG, 88  
Brooks P, 65  
Brott BC, 2  
Brown JS, 71  
Brown MJ, 6  
Brown RD Jr, 31  
Bucher HC, 4  
Bucknall CE, 94  
Byrne GJ, 69  
Bytzer P, 42
- Cadarette SM, 37  
Camargo CA Jr, 46  
Camilleri M, 19
- Campanella SG, 36  
Cannon CP, 49  
CARE-COAD1 Group, 29  
Carter Y, 100  
Casey D, 108  
Castaigne A, 6  
Champs Study Group, 81  
Chandrasekar PH, 59  
Chapman GD 51  
Chea-Woo E, 110  
Childhood Asthma Management Program Research Group, 99  
Chouinard E, 25  
Christens P, 33  
Church TR, 103  
Cochrane Tobacco Addiction Review Group, 60  
Cohen G, 95  
Collins S, 58  
Colly LP, 62  
Colorectal Meta-analysis Collaboration, 25  
Conway DL, 87  
Conway DSG, 52  
Cook DJ, 1  
Corle D, 19  
Coster S, 89  
Cotton MM, 94  
Craig JC, 26  
Cromheecke ME, 62  
Crotty M, 65  
Cryer C, 100  
Cutler BS, 105
- D'Arcy CA, 115  
Dagg KD, 94  
Datta M, 110  
Dawson DL, 105  
De Bruyn G, 57  
de Virgilio C, 117  
Deeks J, 43  
Deeks JJ, 29  
Del Mar C, 109  
Delclaux C, 93  
Deurenberg P, 67  
Diercks DB, 47  
Dimmock PW, 83  
Dirschinger J, 50  
Dixon AS, 106  
Djulbegovic B, 76  
Donnan GA, 12  
Donohue L, 86  
Donovan S, 100  
Douketis JD, 72  
Douwes J, 32  
Ducharme FM, 99
- Early Breast Cancer Trialists' Collaborative Group, 66  
Easton V, 113  
Ebell MH, 28  
Edmonds ML, 46  
Eggenberger ER, 12
- Eikelboom JW, 3  
El Hajjam M, 114  
Elliott CG, 114  
Elliott MW, 15  
Enkin M, 86  
Ernst AA, 47  
Ernst E, 96  
Ettinger M, 64  
European Cancer Prevention Organisation Study Group, 104  
Evans J, 11  
Evans M, A-16 (No. 2)
- Fahey T, 9  
Faivre J, 104  
Feagan B, 20-21  
Feder G, 100  
Feldman W, 108  
Fields SD, 71, 100  
Fink HA, 30  
Fitzgerald JT, 59  
Fitzmaurice DA, 73  
Fletcher RH, 104  
Flewelling D, 28  
Flynn CA, 28  
Flynn PJ, 24  
Frank RM, 69  
Freemantle N, 84  
FRISC II Investigators, 2
- Giacomini MK, A-11-12 (No. 1)  
Gibson NE, 57  
Gillum LA, 72  
Glasziou P, 109  
Glatt AE, 102  
Goldberg AS, 107  
Goldbourr U, 48  
Gonzales O, 87  
Goodman S, 5  
Goodwin PJ, 56  
Griffin JJ, 5  
Grossman E, 48  
Gudmundsson S, 70  
Guidelines' Development Group, 100  
Gulliford MC, 89  
Gunnarsson R, 111  
Gutierrez M, 110  
Guyatt GH, 4
- Haas SB, 54  
Hadenos G, 8  
Hahn S, 57  
Haidar J, 67  
Haines T, 115  
Hamdan MH, 52  
Hankey GJ, 70  
Hannah ME, 112  
Hannah WJ, 112  
Hansson L, 7  
Hardern R, A-16 (No. 1)  
Harkless LB, 107  
Hautvast JG, 67

- Havranek EP, 92  
Hawkins RA, 111  
Haydel MJ, 74  
Haynes RB, A-11-13 (No. 2)  
Hedner T, 7  
Helgason S, 70  
Hengstler P, 4  
Hewson SA, 112  
Hiatt WR, 27, 105  
Hillegass WB, 14, 51  
Hinchey JA, 31  
Hirsh J, 13  
Hobbs FD, 73  
Hodson EM, 26  
Hoffman JR, 75  
Hoffman RM, 91  
Holloway RG, 31  
Homel P, 41  
Honig PK, 17, 96  
Hotopf M, 84  
Hughes CM, 93  
Hunink MG, 27  
Hunt DL, 9  
Huntley A, 96  
Husted S, 2  
Hutchinson A, 68  
  
Igoumenidis NE, 53  
Imperiale TF, 18  
IOEZ Study Group, 88  
Iorio A, 55  
  
Jacobs JW, 85  
Jacobs LD, 81  
Jaglal SB, 37  
Jailwala J, 18  
Joglar JA, 52  
Johnston SC, 72  
Jones G, 65  
Jones GW, 66  
Jones PW, 83  
Josefsson AM, 34, 35  
Jull AB, 14  
  
Kastrati A, 50  
Kaufman DI, 12  
Kelder JC, 51  
Kellett MA, 1  
Kendrick A, 84  
Khurana V, 18-19  
Kiel RJ, 45  
Kirch W, 82  
Klassen T, 36  
Klastersky J, 76  
Klein SR, 117  
Knight JF, 26  
Koch R, 82  
Kochiadakis GE, 53  
Kong S, 19  
Korthals-de Bos IB, 116  
Kramer L, 106  
Kreiger N, 37  
  
Kroenke K, 18  
Kronborg O, 104  
  
L'Her E, 93  
Lagerqvist B, 2  
Lancaster T, 60  
Langer O, 87  
Lanza E, 19  
Laskey WK, 3  
Lassen AT, 42  
Latchaw RE, 8  
Lau JY, 44  
Laville M, 41  
Lawlor PG, 69  
Lederle FA, 30  
Lee KK, 44  
Leversha AM, 36  
Levi M, 62  
Levinson PD, 61  
Lichter EA, 22  
Lifshitz E, 106  
Lindgärde, F, 61  
Little P, 109  
Little P, 59  
Locke KA, 116  
Loder E, 58  
Loprinzi CL, 24  
Luckmann R, 60  
Lumley J, 86  
Lund-Johansen P, 7  
  
MacIntosh C, 9  
Magnusson PK, 34  
Malmberg K, 3  
Mamidipudi SK, 72  
Mandel JS, 103  
Margois DJ, 77  
Marketou ME, 53  
Martinez ME, 21  
Massel D, 50  
McAlister FA, 29  
McDonald AC, 1  
McGee S, 115  
McGeer AJ, 117  
McIntosh A, 68  
McIvor RA, 15  
McQuay H, 58  
Melbye M, 33  
Melnikow J, 34-35  
Meltzer MI, 90  
Messerli FH, 44  
Mesurole B, 118  
Michaels JA, 105  
Milch C, 28  
Mills TJ, 74  
Moayyedi P, 43  
Montgomery AA, 9  
Montori CM, 68  
Montori V, 48, 88, 89  
Morgan DA, 108  
Moride Y, 10  
Morrison LJ, 1  
  
Mortimer PS, 56  
Mosca P, 54  
Mower WR, 75  
Mulrow CD, 6-7, 92  
Muñoz S, 65  
Munro AJ, 23  
Murray ET, 73  
  
National Emergency X-Radiography  
Utilization Study Group, 75  
Nelson M, 11  
Nelson, EA, 77  
Niewwoehner DE, 98  
Nikodem C, 112  
NORDIL Study Group, 7  
Northcutt AR, 19  
Nuovo J, 34-35  
Nybo Andersen AM, 33  
  
O'Brien PM, 83  
O'Malley PG, 85  
Olsen J, 33  
OPUS-1 Investigators, 5  
Orlistat Swedish Multimorbidity  
Study Group, 61  
Orwoll E, 64  
Owen JL, 15  
  
Paesmans M, 76  
Page J, A-11-12 (No. 3)  
Palmer CR, 6  
Pandya KJ, 24  
Parameswaran K, 97  
Park S, 47  
Paty D, 82  
Paulseth JR, 63  
Peacock JL, 56  
Pearce J, 10  
Pearce N, 32  
Pedersen FM, 42  
Perrault A, 10  
Peters J, 46  
Peters J, 68  
Peters TJ, 9  
Peterson M, 54  
Peterson W, 44  
Petrovic M, 108  
Petty GW, 31  
Pettersson G, 70  
Phelps KR, 64  
Phillips C, 92  
Phoenix Colon Cancer Prevention  
Physicians' Network, 21  
Pignone M, 92  
Plant PK, 15  
Polyp Prevention Trial Group, 20  
Potter A, 86  
Powrie JK, 89  
Prasad AS, 59  
Preston CA, 74  
Prostate Cancer Trialists'  
Collaborative Group, 23

- Pulmonary Embolism Prevention (PEP) Trial Collaborative Group, 13
- Qanadli SD, 114
- Quin G, A-15 (No. 3)**
- Ramaker C, 63
- Ramaswamy K, 52
- Räth U, 104
- Raubertas RF, 24
- Rees PJ, 94-95**
- Reginster JY, 37**
- Reisman MA, 5
- Reyzelman AM, 107
- Rich MM, 103**
- Richer B, 17
- Riedemann P, 65
- Roberts RJ, 108
- Roe DJ, 21
- Rogers AI, 18-19**
- Rogers P, 45
- Ronco C, 41
- Rosenfeld M, 111
- Rostom A, 101
- Roth CS, 30
- Rowe BH, 46, 97
- Rubenstein EB, 76
- Ruffin R, 32**
- Ryan EA, 87**
- Sackett DL, 29
- Salazar-Lindo E, 110**
- Santisteban-Ponce J, 110
- Saunders LD, 46
- Sawadsky BV, 1
- Schaffalitzky de Muckadell OB, 42
- Schatzkin A, 20
- Schindler C, 4
- Schmitt BP, 73
- Schömig A, 50**
- Schwartz MD, 43**
- Schweizer J, 82
- Seed PT, 89
- Sharp DJ, 9
- Sigurdsson JA, 70
- Silagy C, 60
- Simmer K, 67**
- Simon JH, 81
- Singh N, 45
- Skwarska E, 95
- Skwarski KM, 95
- Small R, 86
- Smeeth L, 11
- Smetana S, 91
- Smith BJ, 29, 32**
- Smith RC, 83**
- Soo S, 43
- Sørensen P, 35
- Sowden A, 60
- Stead L, 60
- Steele BT, 26**
- Stent versus Thrombolysis for Occluded Coronary Arteries in Patients with Acute Myocardial Infarction Study Investigators, 50
- Straus SE, 29
- Strom BL, 77
- Stuart CA, 88
- Study of Osteoporotic Fractures Research Group, 71
- Study Section on Infections of Multinational Association for Supportive Care in Cancer, 76
- Sung JJ, 44
- Suttorp MJ, 51
- Swaminathan R, 89
- Takakuwa KM, 47
- Talley NJ, 101**
- Taylor PR, 30**
- ten Berg JM, 51
- Term Breech Trial Collaborative Group, 112
- Thomas M, 109**
- Thomas SM, 113**
- Thompson WW, 90
- Todd KH, 75
- Tomkins G, 85
- Topol EJ, 49
- Trobe JD, 12
- Trombello KA, 107
- Tugwell P, 101
- Umeta M, 67
- van Dielen, 116
- van Hilten JJ, 63
- van Tulder MW, 116
- Vayser DH, 107
- Veldhuyzen van Zanten SJO, 42**
- Verbeek PR, 1
- Visser K, 27
- Virthinghoff E, 71
- The Voluntary HIV-1 Counseling and Testing Efficacy Study Group, 22
- Wagener MM, 45
- Waldenström U, 86
- Wall DB, 117
- Wallentin L, 2
- Walters EH, 98
- Walters J, 98
- Wardlaw JM, 113
- Waters J, 14
- Weaver WD, 5
- Weinstein T, 91
- Weiss S, 64
- Weiss SJ, 47
- Welss G, 101
- West CE, 67
- Westrich GH, 54**
- Whisnant JP, 31
- Whitaker JN, 12
- White PM, 113
- Wiffen P, 58
- Wilkinson D, 102
- Willis NS, 26
- Wilson MW, A-15 (No. 1)**
- Wohlfahrt J, 33
- Wolfson AB, 75
- Wolfson C, 10
- Wormald R, 11
- Wyatt KM, 83
- Wyer P, 47**
- Wyman JF, 71
- Xenakis EM-J, 87
- Yarnall SR, 53**
- Ylitalo N, 34, 35
- Yu VL, 45
- Zaroukian MH, A-15 (No. 2)**
- Zucker MI, 75

## A

- Abortion, spontaneous
  - Higher maternal age was associated with increased risks for fetal death and ectopic pregnancy, 33
- Accidental falls
  - Weekly urge urinary incontinence was associated with increased risk for falls and nonspinal fractures in older women, 71
- Adenoma
  - A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20
  - A high-fiber cereal supplement did not prevent the recurrence of colorectal adenomas, 21
- Adrenal cortex hormones
  - Review: Corticosteroids do not improve visual acuity in acute monosymptomatic optic neuritis over the long term, 12
- Adrenergic beta-agonists
  - Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at preventing cardiovascular mortality and morbidity, 7
  - On-demand use of  $\beta_2$ -agonists led to better asthma control than did regular use in moderate-to-severe asthma, 17
- Albuterol
  - On-demand use of  $\beta_2$ -agonists led to better asthma control than did regular use in moderate-to-severe asthma, 17
  - Spacers were better and less expensive than nebulizers for giving albuterol to children with moderate-to-severe acute asthma, 36
- Alendronate
  - Alendronate increased bone mineral density and decreased vertebral fractures in men with osteoporosis, 64
- Allergens
  - Review: Evidence does not support an association between allergen levels and risk for asthma, 32
- Alzheimer disease
  - Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10
- Ambulatory care
  - Self-management of long-term oral anticoagulation was as effective as specialist anticoagulation clinic management, 62
- Amiloride
  - Long-acting nifedipine was as effective as hydrochlorothiazide plus amiloride for reducing mortality and morbidity in hypertension, 6
- Amiodarone
  - Amiodarone maintained sinus rhythm better than did sotalol or placebo in atrial fibrillation, but adverse effects were more frequent, 53
- Androgen antagonists
  - Review: Maximum androgen blockade does not confer additional survival benefit to androgen suppression in prostate cancer, 23
- Angina
  - Review: Glycoprotein IIb/IIIa inhibitors reduce combined end points in acute coronary syndromes, 49
  - Review: PTCA is associated with less angina but more CABG in patients with nonacute CAD than is medical treatment, 4
- Angioplasty, transluminal percutaneous, coronary
  - Coumarins and aspirin reduced clinical events more than did aspirin alone when drugs were started before scheduled PTCA, 51
  - An invasive strategy reduced death, myocardial infarction, and readmissions in unstable coronary artery disease, 2
  - Review: PTCA is associated with less angina but more CABG in patients with nonacute CAD than is medical treatment, 4
  - Routine stenting improved outcomes and had similar costs to that of balloon angioplasty with provisional stenting, 5
- Angiotensin-converting enzyme inhibitors
  - Review: Intensive blood-pressure control and drugs reduce morbidity and mortality in hypertension and diabetes mellitus, 48
- Antacids
  - Review: Prokinetics and histamine-2 receptor antagonists improve symptom scores in nonulcer dyspepsia, 43
- Anti-arrhythmia agents
  - Amiodarone maintained sinus rhythm better than did sotalol or placebo in atrial fibrillation, but adverse effects were more frequent, 53
- Antibiotics
  - Antibiotic monotherapy with possible stopping at 3 days benefited low-risk ICU patients with pulmonary infiltrates, 45
  - Review: Antibiotics reduce the duration and severity of travelers' diarrhea, 57
- Anticoagulants
  - Coumarins and aspirin reduced clinical events more than did aspirin alone when drugs were started before scheduled PTCA, 51
  - An invasive strategy reduced death, myocardial infarction, and readmissions in unstable coronary artery disease, 2
  - A nurse-led clinic and computer decision-support software for anticoagulation decisions were as effective as a hospital clinic, 73
  - Review: Pneumatic compression and LMWH are more effective for preventing DVT after total knee arthroplasty than is aspirin or warfarin, 54
  - Self-management of long-term oral anticoagulation was as effective as specialist anticoagulation clinic management, 62
- Anticonvulsants
  - Review: Anticonvulsant drugs relieve chronic but not acute pain, 58
- Antihypertensive agents
  - Review: Intensive blood-pressure control and drugs reduce morbidity and mortality in hypertension and diabetes mellitus, 48
- Anti-inflammatory agents
  - Review: Corticosteroids do not improve visual acuity in acute monosymptomatic optic neuritis over the long term, 12
- Antineoplastic agents
  - A clinical score identified cancer patients with febrile neutropenia at low risk for complications, 76
  - Review: Palliative chemotherapy reduces death and progression at 12 months in advanced or metastatic colorectal cancer, 25
- Aortic aneurysm, abdominal
  - A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20
- Arthritis, psoriatic
  - Review: Sulfasalazine, azathioprine, and etretinate improve a global index of disease activity in psoriatic arthritis, 65
- Aspirin
  - Low-dose aspirin reduced DVT and pulmonary embolism but increased postoperative bleeding requiring transfusion in hip-surgery patients, 13
  - Review: Pneumatic compression and LMWH are more effective for preventing DVT after total knee arthroplasty than is aspirin or warfarin, 54
- Asthma
  - On-demand use of  $\beta_2$ -agonists led to better asthma control than did regular use in moderate-to-severe asthma, 17
  - Review: Evidence does not support an association between allergen levels and risk for asthma, 32

- Review: Inhaled corticosteroids alone appear to be as effective as oral corticosteroids after ED discharge for acute asthma, 46  
 Spacers were better and less expensive than nebulizers for giving albuterol to children with moderate-to-severe acute asthma, 36
- Athroplasty, replacement, knee  
 Review: Pneumatic compression and LMWH are more effective for preventing DVT after total knee athroplasty than is aspirin or warfarin, 54
- Atrial fibrillation  
 Amiodarone maintained sinus rhythm better than did sotalol or placebo in atrial fibrillation, but adverse effects were more frequent, 53  
 An initial energy of 360 J for elective cardioversion was more effective than 100 or 200 J in persistent atrial fibrillation, 52
- Azathioprine  
 Review: Sulfasalazine, azathioprine, and etretinate improve a global index of disease activity in psoriatic arthritis, 65

## B

- Bandages  
 Multilayer bandaging plus compression hosiery was better than hosiery alone for unilateral lymphedema of a limb, 56  
 Review: Pentoxifylline alone or as an adjunct to compression therapy improves healing in venous leg ulcers, 14  
 Review: Pneumatic compression and LMWH are more effective for preventing DVT after total knee athroplasty than is aspirin or warfarin, 54  
 A simple model based on wound size and duration predicted healing of venous leg ulcers at 24 weeks, 77
- Behavior therapy  
 Review: Various interventions increase smoking cessation rates, 60
- Bone density  
 Alendronate increased bone mineral density and decreased vertebral fractures in men with osteoporosis, 64
- Brain ischemia  
 Review: Current oral contraceptive use increases the risk for ischemic stroke, 72
- Breast neoplasms  
 Clonidine was effective for reducing tamoxifen-associated hot flashes in postmenopausal women with breast cancer, 24  
 Review: Radiotherapy reduces death from and recurrence of breast cancer but increases death from other causes, 66
- Bromocriptine  
 Review: Bromocriptine may be beneficial in delaying motor complications and dyskinesias in Parkinson disease, 63

## C

- Calcium channel blockers  
 Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at preventing cardiovascular mortality and morbidity, 7  
 Long-acting nifedipine was as effective as hydrochlorothiazide plus amiloride for reducing mortality and morbidity in hypertension, 6  
 Review: Intensive blood-pressure control and drugs reduce morbidity and mortality in hypertension and diabetes mellitus, 48

- Cardiovascular diseases  
 Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at preventing cardiovascular mortality and morbidity, 7
- Cereals  
 A high-fiber cereal supplement did not prevent the recurrence of colorectal adenomas, 21
- Cerebrovascular accident  
 Ischemic stroke subtypes had different short- and long-term functional outcomes, mortality, and recurrence rates, 31  
 Review: Current oral contraceptive use increases the risk for ischemic stroke, 72  
 Review: Stroke centers need administrative support, protocols, multidisciplinary staff, and access to hospital services, 8
- Cervical neoplasms  
 A higher level of human papillomavirus 16 DNA was associated with an increased risk for cervical carcinoma in situ, 34  
 Persistently high loads of human papillomavirus 16 over time were associated with an increased risk for cervical cancer, 35
- Cervical vertebrae  
 Absence of 5 clinical criteria ruled out cervical-spine injury in blunt trauma in the emergency department, 75
- Chronic disease  
 Review: Anticonvulsant drugs relieve chronic but not acute pain, 58
- Ciprofloxacin  
 Antibiotic monotherapy with possible stopping at 3 days benefited low-risk ICU patients with pulmonary infiltrates, 45
- Cisapride  
 Review: Prokinetics and histamine-2 receptor antagonists improve symptom scores in nonulcer dyspepsia, 43
- Colonic diseases, functional  
 Alosetron was effective and safe for relieving abdominal symptoms in women with the irritable bowel syndrome, 19  
 Review: Smooth-muscle relaxants treat abdominal pain and loperamide reduces diarrhea in the irritable bowel syndrome, 18
- Colorectal neoplasms  
 A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20  
 A high-fiber cereal supplement did not prevent the recurrence of colorectal adenomas, 21  
 Review: Palliative chemotherapy reduces death and progression at 12 months in advanced or metastatic colorectal cancer, 25
- Common cold  
 Zinc acetate lozenges reduced the duration and severity of symptoms of the common cold, 59
- Consumer participation  
*Smart Health Choices: How to Make Informed Health Decisions* (Resource Corner), A-16 (No. 2)
- Contraceptives, oral  
 Review: Current oral contraceptive use increases the risk for ischemic stroke, 72
- Coronary artery bypass  
 Review: PTCA is associated with less angina but more CABG in patients with nonacute CAD than is medical treatment, 4
- Coronary disease  
 Coumarins and aspirin reduced clinical events more than did aspirin alone when drugs were started before scheduled PTCA, 51  
 An invasive strategy reduced death, myocardial infarction, and readmissions in unstable coronary artery disease, 2



- Orlistat with diet was effective and safe for weight loss and coronary risk reduction in obesity, 61
- Review: No convincing difference exists between unfractionated and low-molecular-weight heparin in the acute coronary syndrome, 3
- Routine stenting improved outcomes and had similar costs to that of balloon angioplasty with provisional stenting, 5
- Corticosteroids
  - Review: Inhaled corticosteroids alone appear to be as effective as oral corticosteroids after ED discharge for acute asthma, 46
- Cough
  - Zinc acetate lozenges reduced the duration and severity of symptoms of the common cold, 59
- Coumarins
  - Coumarins and aspirin reduced clinical events more than did aspirin alone when drugs were started before scheduled PTCA, 51
- Counseling
  - Voluntary counseling and testing reduced unprotected intercourse among adults in 3 developing countries, 22
- Cranio-cerebral trauma
  - A decision rule with 7 clinical criteria identified patients with minor head injury who did not need computed tomography, 74

## D

- Databases, bibliographic
  - PubMed clinical queries: a Web tool for filtered retrieval of citations relevant to evidence-based practice (Resource Corner), A-15 (No. 2)
- Decision support techniques
  - A nurse-led clinic and computer decision-support software for anticoagulation decisions were as effective as a hospital clinic, 73
- Decision-making, computer-assisted
  - A computer-based decision support system and risk chart did not reduce cardiovascular risk or blood pressure in hypertension, 9
- Dementia
  - The Hopkins Verbal Learning Test had high sensitivity and good specificity for detecting mild dementia in older persons, 69
- Densitometry
  - A simple clinical tool that included age, weight, and estrogen use helped to select women for bone densitometry, 37
- Diabetes mellitus
  - Orlistat with diet was effective and safe for weight loss and coronary risk reduction in obesity, 61
  - Review: Intensive blood-pressure control and drugs reduce morbidity and mortality in hypertension and diabetes mellitus, 48
- Diabetic retinopathy
  - Review: Mydriatic retinal photography is the most effective test for detecting diabetic retinopathy, 68
- Diagnostic techniques and procedures
  - Review: Mydriatic retinal photography is the most effective test for detecting diabetic retinopathy, 68
- Diarrhea
  - Review: Antibiotics reduce the duration and severity of travelers' diarrhea, 57
- Diet, fat-restricted
  - A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20
- Dietary fiber
  - A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20

- A high-fiber cereal supplement did not prevent the recurrence of colorectal adenomas, 21
- Dietary supplements
  - Zinc supplements increased growth more in stunted infants than in nonstunted infants, 67
- Diltiazem
  - Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at preventing cardiovascular mortality and morbidity, 7
- Diuretics
  - Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at preventing cardiovascular mortality and morbidity, 7
  - Long-acting nifedipine was as effective as hydrochlorothiazide plus amiloride for reducing mortality and morbidity in hypertension, 6
- Donepezil
  - Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10
- Drug monitoring
  - Self-management of long-term oral anticoagulation was as effective as specialist anticoagulation clinic management, 62
- Drug therapy
  - Orlistat with diet was effective and safe for weight loss and coronary risk reduction in obesity, 61
- Dyspepsia
  - Helicobacter pylori* testing plus eradication was as effective and safe as prompt endoscopy for dyspepsia, 42
  - Review: Prokinetics and histamine-2 receptor antagonists improve symptom scores in nonulcer dyspepsia, 43

## E

- Electric countershock
  - An initial energy of 360 J for elective cardioversion was more effective than 100 or 200 J in persistent atrial fibrillation, 52
- Emergency medicine
  - Best BETS (Resource Corner), A-15 (No. 3)
- Endoscopy
  - Helicobacter pylori* testing plus eradication was as effective and safe as prompt endoscopy for dyspepsia, 42
  - A high-dose infusion of omeprazole after endoscopic treatment of bleeding peptic ulcers reduced recurrent bleeding, 44
- Epinephrine
  - A high-dose infusion of omeprazole after endoscopic treatment of bleeding peptic ulcers reduced recurrent bleeding, 44
- Estrogen
  - Review: Current oral contraceptive use increases the risk for ischemic stroke, 72
- Evidence-based medicine
  - Evidence-Based Clinical Practice* (Resource Corner), A-16 (No. 1)
  - Evidence-Based Medicine: How to Practice and Teach EBM*, (Resource Corner), A-15 (No. 1)
  - Of studies, syntheses, synopses, and systems: the "4S" evolution of services for finding current best evidence (Editorial), A-11-13 (No. 2)
  - PubMed clinical queries: a Web tool for filtered retrieval of citations relevant to evidence-based practice (Resource Corner), A-15 (No. 2)
  - The rocky road: qualitative research as evidence (Editorial), A-11-13 (No. 1)

*Smart Health Choices: How to Make Informed Health Decisions*  
(Resource Corner), A-16 (No. 2)  
Using Evidence in Health and Social Care (Resource  
Corner), A-16 (No. 3)

## F

- Fenoterol  
On-demand use of  $\beta_2$ -agonists led to better asthma control than did  
regular use in moderate-to-severe asthma, 17
- Fetal death  
Higher maternal age was associated with increased risks for fetal  
death and ectopic pregnancy, 33
- Fever  
A clinical score identified cancer patients with febrile neutropenia  
at low risk for complications, 76
- Fractures, spontaneous  
Weekly urge urinary incontinence was associated with increased risk  
for falls and nonspinal fractures in older women, 71

## G

- Gadolinium  
Review: Gadolinium-enhanced magnetic resonance angiography is  
highly sensitive and specific for peripheral arterial disease, 27
- Gastrointestinal transit  
Alosetron was effective and safe for relieving abdominal symptoms in  
women with the irritable bowel syndrome, 19  
Review: Smooth-muscle relaxants treat abdominal pain and  
loperamide reduces diarrhea in the irritable bowel syndrome, 18
- Ginkgo biloba*  
Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are  
more effective than placebo in Alzheimer disease, 10
- Growth disorders  
Zinc supplements increased growth more in stunted infants than  
in nonstunted infants, 67

## H

- Health care costs  
Routine stenting improved outcomes and had similar costs to that of  
balloon angioplasty with provisional stenting, 5
- Health promotion  
Review: Various interventions increase smoking cessation rates, 60
- Helicobacter pylori*  
*Helicobacter pylori* testing plus eradication was as effective and safe as  
prompt endoscopy for dyspepsia, 42
- Hemofiltration  
Continuous hemofiltration at an ultrafiltration rate  $\geq 35$  mL/h per  
kg of body weight improved survival in acute renal failure, 41
- Heparin  
Review: Heparin reduces venous thromboembolism in neurosurgery  
but increases the risk for bleeding, 55  
Review: No convincing difference exists between unfractionated and  
low-molecular-weight heparin in the acute coronary syndrome, 3
- Herpes zoster  
Postherpetic neuralgia was not frequent or severe after a first episode  
of herpes zoster, 70
- Histamine  $H_2$  antagonists  
Review: Prokinetics and histamine-2 receptor antagonists improve  
symptom scores in nonulcer dyspepsia, 43

- HIV-1  
Voluntary counseling and testing reduced unprotected intercourse  
among adults in 3 developing countries, 22
- Hospitals, special  
Review: Stroke centers need administrative support, protocols,  
multidisciplinary staff, and access to hospital services, 8
- Hot flashes  
Clonidine was effective for reducing tamoxifen-associated hot flashes  
in postmenopausal women with breast cancer, 24
- Hydrochlorothiazide  
Long-acting nifedipine was as effective as hydrochlorothiazide  
plus amiloride for reducing mortality and morbidity in  
hypertension, 6
- Hypertension  
A computer-based decision support system and risk chart did not  
reduce cardiovascular risk or blood pressure in hypertension, 9  
Diltiazem was as effective as diuretics or  $\beta$ -blockers, or both, at  
preventing cardiovascular mortality and morbidity, 7  
Long-acting nifedipine was as effective as hydrochlorothiazide  
plus amiloride for reducing mortality and morbidity in  
hypertension, 6  
Review: Intensive blood-pressure control and drugs reduce morbidity  
and mortality in hypertension and diabetes mellitus, 48

## I

- Information services  
Of studies, syntheses, synopses, and systems: the "4S" evolution of  
services for finding current best evidence (Editorial), A-11-13  
(No. 2)
- Intensive care units  
Antibiotic monotherapy with possible stopping at 3 days benefited  
low-risk ICU patients with pulmonary infiltrates, 45
- Intubation  
Noninvasive ventilation reduced deaths and the need for intubation  
in acute-on-chronic obstructive pulmonary disease, 15

## K

- Kidney failure, acute  
Continuous hemofiltration at an ultrafiltration rate  $\geq 35$  mL/h per  
kg of body weight improved survival in acute renal failure, 41

## L

- Lecithin  
Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba*  
are more effective than placebo in Alzheimer disease, 10
- Leg ulcers  
Review: Pentoxifylline alone or as an adjunct to compression therapy  
improves healing in venous leg ulcers, 14
- Levodopa  
Review: Bromocriptine may be beneficial in delaying motor compli-  
cations and dyskinesias in Parkinson disease, 63
- Linopirdine  
Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are  
more effective than placebo in Alzheimer disease, 10
- Lung diseases, obstructive  
Noninvasive ventilation reduced deaths and the need for intubation  
in acute-on-chronic obstructive pulmonary disease, 15  
Smoking, age, and laryngeal height predicted obstructive airway  
disease in adults, 29

Lymphedema

- Multilayer bandaging plus compression hosiery was better than hosiery alone for unilateral lymphedema of a limb, 56

M

Macular degeneration

- Review: Photodynamic therapy prevents vision loss in neovascular age-related macular degeneration, 11

Magnetic resonance angiography

- Review: Gadolinium-enhanced magnetic resonance angiography is highly sensitive and specific for peripheral arterial disease, 27

Mass screening

- The Hopkins Verbal Learning Test had high sensitivity and good specificity for detecting mild dementia in older persons, 69

Medline

- PubMed clinical queries: a Web tool for filtered retrieval of citations relevant to evidence-based practice (Resource Corner), A-15 (No. 2)

Metrifonate

- Review: Donepezil, metrifonate, rivastigmine, and Ginkgo biloba are more effective than placebo in Alzheimer disease, 10

Models, statistical

- A clinical score identified cancer patients with febrile neutropenia at low risk for complications, 76

Mortality

- Mechanical ventilation with a lower tidal volume decreased mortality in the acute respiratory distress syndrome, 16
- Review: No convincing difference exists between unfractionated and low-molecular-weight heparin in the acute coronary syndrome, 3

Myocardial infarction

- Coronary stenting and platelet glycoprotein inhibitors were more effective than was t-PA for acute myocardial infarction, 50
- An invasive strategy reduced death, myocardial infarction, and readmissions in unstable coronary artery disease, 2
- Review: The accuracy of troponin T and I tests in acute MI depends on timing after onset of symptoms, 28
- Review: Glycoprotein IIb/IIIa inhibitors reduce combined end points in acute coronary syndromes, 49
- Review: No convincing difference exists between unfractionated and low-molecular-weight heparin in the acute coronary syndrome, 3
- Review: Prehospital thrombolysis for acute myocardial infarction decreases short-term hospital mortality, 1
- Review: PTCA is associated with less angina but more CABG in patients with nonacute CAD than is medical treatment, 4

N

Nebulizers and vaporizers

- Spacers were better and less expensive than nebulizers for giving albuterol to children with moderate to severe acute asthma, 36

Neoplasia recurrence, local

- A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20
- A high-fiber cereal supplement did not prevent the recurrence of colorectal adenomas, 21

Nephrotic syndrome

- Review: 3 to 7 months of prednisone is more effective than 2 months in children with the steroid-responsive nephrotic syndrome, 26

Neuralgia

- Postherpetic neuralgia was not frequent or severe after a first episode of herpes zoster, 70

Neurology

- Review: Corticosteroids do not improve visual acuity in acute monosymptomatic optic neuritis over the long term, 12

Neurosurgical procedures

- Review: Heparin reduces venous thromboembolism in neurosurgery but increases the risk for bleeding, 55

Neutropenia

- A clinical score identified cancer patients with febrile neutropenia at low risk for complications, 76

Nifedipine

- Long-acting nifedipine was as effective as hydrochlorothiazide plus amiloride for reducing mortality and morbidity in hypertension, 6

Nurse practitioners

- A nurse-led clinic and computer decision-support software for anticoagulation decisions were as effective as a hospital clinic, 73

O

Obesity

- Orlistat with diet was effective and safe for weight loss and coronary risk reduction in obesity, 61

Omeprazole

- A high-dose infusion of omeprazole after endoscopic treatment of bleeding peptic ulcers reduced recurrent bleeding, 44

Optic neuritis

- Review: Corticosteroids do not improve visual acuity in acute monosymptomatic optic neuritis over the long term, 12

Osteoporosis

- Alendronate increased bone mineral density and decreased vertebral fractures in men with osteoporosis, 64
- A simple clinical tool that included age, weight, and estrogen use helped to select women for bone densitometry, 37

P

Pain

- Postherpetic neuralgia was not frequent or severe after a first episode of herpes zoster, 70
- Review: Anticonvulsant drugs relieve chronic but not acute pain, 58

Palliative care

- Review: Palliative chemotherapy reduces death and progression at 12 months in advanced or metastatic colorectal cancer, 25

Palpation

- A diet low in fat and high in fiber, fruits, and vegetables did not reduce the recurrence of colorectal adenomas, 20

Papillomavirus, human

- A higher level of human papillomavirus 16 DNA was associated with an increased risk for cervical carcinoma in situ, 34
- Persistently high loads of human papillomavirus 16 over time were associated with an increased risk for cervical cancer, 35

Parkinson disease

- Review: Bromocriptine may be beneficial in delaying motor complications and dyskinesias in Parkinson disease, 63

Patient selection

- A simple clinical tool that included age, weight, and estrogen use helped to select women for bone densitometry, 37



## Pentoxifylline

Review: Pentoxifylline alone or as an adjunct to compression therapy improves healing in venous leg ulcers, 14

## Peptic ulcer hemorrhage

A high-dose infusion of omeprazole after endoscopic treatment of bleeding peptic ulcers reduced recurrent bleeding, 44

## Peptic ulcer

A high-dose infusion of omeprazole after endoscopic treatment of bleeding peptic ulcers reduced recurrent bleeding, 44

## Peripheral vascular diseases

Review: Gadolinium-enhanced magnetic resonance angiography is highly sensitive and specific for peripheral arterial disease, 27

## Phototherapy

Review: Photodynamic therapy prevents vision loss in neovascular age-related macular degeneration, 11

## Physical examination

Smoking, age, and laryngeal height predicted obstructive airway disease in adults, 29

## Pirenzepine

Review: Prokinetics and histamine-2 receptor antagonists improve symptom scores in nonulcer dyspepsia, 43

## Platelet aggregation inhibitors

Low-dose aspirin reduced DVT and pulmonary embolism but increased postoperative bleeding requiring transfusion in hip-surgery patients, 13

Review: Glycoprotein IIb/IIIa inhibitors reduce combined end points in acute coronary syndromes, 49

## Platelet glycoprotein GPIIb/IIIa complex

Coronary stenting and platelet glycoprotein inhibitors were more effective than was t-PA for acute myocardial infarction, 50

Review: Glycoprotein IIb/IIIa inhibitors reduce combined end points in acute coronary syndromes, 49

## Pneumonia

Antibiotic monotherapy with possible stopping at 3 days benefited low-risk ICU patients with pulmonary infiltrates, 45

## Postoperative complications

Coumarins and aspirin reduced clinical events more than did aspirin alone when drugs were started before scheduled PTCA, 51

Low-dose aspirin reduced DVT and pulmonary embolism but increased postoperative bleeding requiring transfusion in hip-surgery patients, 13

## Prednisone

Review: 3 to 7 months of prednisone is more effective than 2 months in children with the steroid-responsive nephrotic syndrome, 26

## Pregnancy

Higher maternal age was associated with increased risks for fetal death and ectopic pregnancy, 33

## Prochlorperazine

Prochlorperazine was more effective than promethazine in uncomplicated nausea and vomiting in the emergency department, 47

## Progesterone

Review: Current oral contraceptive use increases the risk for ischemic stroke, 72

## Promethazine

Prochlorperazine was more effective than promethazine in uncomplicated nausea and vomiting in the emergency department, 47

## Propentofylline

Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10

## Prostatic neoplasms

Review: Maximum androgen blockade does not confer additional survival benefit to androgen suppression in prostate cancer, 23

## Pulmonary embolism

Low-dose aspirin reduced DVT and pulmonary embolism but increased postoperative bleeding requiring transfusion in hip-surgery patients, 13

## Q

## Qualitative research

The rocky road: qualitative research as evidence (Editorial), A-11-13 (No. 1)

## R

## Radiography

A decision rule with 7 clinical criteria identified patients with minor head injury who did not need computed tomography, 74

A simple clinical tool that included age, weight, and estrogen use helped to select women for bone densitometry, 37

## Radiotherapy

Review: Radiotherapy reduces death from and recurrence of breast cancer but increases death from other causes, 66

## Randomized controlled trials

A graphic framework for teaching critical appraisal of randomized controlled trials (Editorial), A-11-12 (No. 3)

## Reading

Of studies, syntheses, synopses, and systems: the "4S" evolution of services for finding current best evidence (Editorial), A-11-13 (No. 2)

## Recurrence

Ischemic stroke subtypes had different short- and long-term functional outcomes, mortality, and recurrence rates, 31

## Respiratory distress syndrome

Mechanical ventilation with a lower tidal volume decreased mortality in the acute respiratory distress syndrome, 16

## Risk assessment

A computer-based decision support system and risk chart did not reduce cardiovascular risk or blood pressure in hypertension, 9

## Rivastigmine

Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10

## S

## Selegiline

Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10

## Self-administration

Self-management of long-term oral anticoagulation was as effective as specialist anticoagulation clinic management, 62

## Sensitivity and specificity

The Hopkins Verbal Learning Test had high sensitivity and good specificity for detecting mild dementia in older persons, 69

Review: Gadolinium-enhanced magnetic resonance angiography is highly sensitive and specific for peripheral arterial disease, 27

- Serotonin antagonists  
 Alosetron was effective and safe for relieving abdominal symptoms in women with the irritable bowel syndrome, 19  
 Review: Smooth-muscle relaxants treat abdominal pain and loperamide reduces diarrhea in the irritable bowel syndrome, 18
- Severity of illness index  
 A clinical score identified cancer patients with febrile neutropenia at low risk for complications, 76
- Sex behavior  
 Voluntary counseling and testing reduced unprotected intercourse among adults in 3 developing countries, 22
- Smoking cessation  
 Review: Various interventions increase smoking cessation rates, 60
- Sotalol  
 Amiodarone maintained sinus rhythm better than did sotalol or placebo in atrial fibrillation, but adverse effects were more frequent, 53
- Spinal injuries  
 Absence of 5 clinical criteria ruled out cervical-spine injury in blunt trauma in the emergency department, 75
- Spirometry  
 Smoking, age, and laryngeal height predicted obstructive airway disease in adults, 29
- Stents  
 Coronary stenting and platelet glycoprotein inhibitors were more effective than was t-PA for acute myocardial infarction, 50
- Sulfasalazine  
 Review: Sulfasalazine, azathioprine, and etretinate improve a global index of disease activity in psoriatic arthritis, 65
- T**
- Tamoxifen  
 Clonidine was effective for reducing tamoxifen-associated hot flashes in postmenopausal women with breast cancer, 24
- Thromboembolism  
 Low-dose aspirin reduced DVT and pulmonary embolism but increased postoperative bleeding requiring transfusion in hip-surgery patients, 13  
 A nurse-led clinic and computer decision-support software for anticoagulation decisions were as effective as a hospital clinic, 73  
 Review: Heparin reduces venous thromboembolism in neurosurgery but increases the risk for bleeding, 55  
 Review: Pneumatic compression and LMWH are more effective for preventing DVT after total knee arthroplasty than is aspirin or warfarin, 54
- Thrombolytic therapy  
 Review: Prehospital thrombolysis for acute myocardial infarction decreases short-term hospital mortality, 1
- Tidal volume  
 Mechanical ventilation with a lower tidal volume decreased mortality in the acute respiratory distress syndrome, 16
- Tissue plasminogen activator  
 Coronary stenting and platelet glycoprotein inhibitors were more effective than was t-PA for acute myocardial infarction, 50
- Tomography, x-ray computed  
 A decision rule with 7 clinical criteria identified patients with minor head injury who did not need computed tomography, 74

- Travel  
 Review: Antibiotics reduce the duration and severity of travelers' diarrhea, 57
- Treatment, emergency  
 Prochlorperazine was more effective than promethazine in uncomplicated nausea and vomiting in the emergency department, 47
- Troponin  
 Review: The accuracy of troponin T and I tests in acute MI depends on timing after onset of symptoms, 28
- Tyrosine  
 Review: Glycoprotein IIb/IIIa inhibitors reduce combined end points in acute coronary syndromes, 49

**U**

- Urinary incontinence  
 Weekly urge urinary incontinence was associated with increased risk for falls and nonspinal fractures in older women, 71

**V**

- Vaginal smears  
 A higher level of human papillomavirus 16 DNA was associated with an increased risk for cervical carcinoma in situ, 34  
 Persistently high loads of human papillomavirus 16 over time were associated with an increased risk for cervical cancer, 35
- Varicose ulcer  
 A simple model based on wound size and duration predicted healing of venous leg ulcers at 24 weeks, 77
- Ventilation  
 Noninvasive ventilation reduced deaths and the need for intubation in acute-on-chronic obstructive pulmonary disease, 15
- Verteporfin  
 Review: Photodynamic therapy prevents vision loss in neovascular age-related macular degeneration, 11
- Vitamin E  
 Review: Donepezil, metrifonate, rivastigmine, and *Ginkgo biloba* are more effective than placebo in Alzheimer disease, 10

**W**

- Weight loss  
 Orlistat with diet was effective and safe for weight loss and coronary risk reduction in obesity, 61
- Wound healing  
 A simple model based on wound size and duration predicted healing of venous leg ulcers at 24 weeks, 77
- Wounds, nonpenetrating  
 Absence of 5 clinical criteria ruled out cervical-spine injury in blunt trauma in the emergency department, 75

**Z**

- Zinc  
 Zinc acetate lozenges reduced the duration and severity of symptoms of the common cold, 59  
 Zinc supplements increased growth more in stunted infants than in nonstunted infants, 67

